



PATIENT NOTES | OLECRANON BURSITIS

A bursa is a fluid filled sack between skin and bone which allows the skin to move easily over the bone. There is a bursa outside the elbow called the olecranon bursa which can be felt around the hard bone of the point of the elbow only with the elbow straight. With the elbow fully bent a normal bursa cannot be felt. The bursa sometimes becomes enlarged as a result of pressure or friction.

The commonest cause of swelling within the bursa, not related to trauma, is gout. There may be a sizeable lump with calcium seen on x-ray. This can also happen in rheumatoid arthritis. Unfortunately it can be difficult to differentiate between an infected and a non infected olecranon bursa and about one in five cases of acute bursitis are, in fact, infected. In general:

- Gradual swelling indicates a chronic or long-lasting condition.
- Sudden swelling indicates a traumatic injury or an infection in the elbow.
- If the elbow was injured, the skin may be scraped or cut.
- Red, hot skin may indicate an infection.
- Pain and tenderness is variable.

Infected bursae are almost always painful whereas non infected bursitis is painful in less than 25% of cases. A swollen olecranon bursa is usually not painful unless it is associated with infection or a specific inflammatory process such as gout or rheumatoid arthritis. In most patients there is no problem with the arm straight, however, symptoms become evident when attempting to bend the elbow beyond 90° or when leaning on the elbow.

Treatment depends on whether the bursa is infected or not. In non infected bursitis the simplest treatment is to stop the elbow moving and allow the fluid filled sac to rest. This often involves a resting splint, a compression bandage and ice packs.

In addition to these measures an infected bursitis will require antibiotic treatment (this may require admission to hospital or simply be in tablet form). In certain situations it may be appropriate to aspirate fluid out of the bursa or possibly to inject a corticosteroid into it. **While this has been shown to reduce the recurrence of the bursa, it also increases the likelihood of infection.** This injection should never be made into an infected olecranon bursa.

For the great majority of patients, simple symptomatic treatment such as using a padded elbow brace or resting the arm on a pillow will be sufficient. Some patients will require repeated periods of immobilisation and some patients will eventually come to surgical removal of the bursa. This procedure is usually extremely effective, however, in a small number of patients the wound does not heal properly for up to three months. A period of immobilisation of the elbow following surgery is necessary and does not guarantee that the bursa will not recur.

In most patients, the swelling is simply a nuisance and as long as the bursa is not infected should not be interfered with.

J. GOLDBERG	shoulder surgery	W. BRUCE	hip, knee	D. SHER	shoulder, elbow, knee
R. PATTINSON	paediatric, general	A. LOEFLER	hip, knee, spine	J. NEGRINE	foot, ankle
A. TURNBULL	hip, knee	P. WALKER	hip, knee	I. POPOFF	knee, shoulder, general
T. GOTHELF	foot, ankle, shoulder, elbow			J. TRANTALIS	shoulder, elbow

SPORTS PHYSICIANS: J.BEST, M.CUSI, P.ANNETT, M.RAFTERY

160 Belmore Road, Randwick NSW 2031
47-49 Burwood Road, Concord NSW 2137
187 Macquarie Street, Sydney 2000
2 Pearl Street, Hurstville 2220

www.orthosports.com.au

Tel: 9399 5333 Fax: 9398 8673
Tel: 9744 2666 Fax: 9744 3706
Tel: 9233 2883 Fax: 9231 0103
Tel: 9580 6066 Fax: 9580 0890

office@orthosports.com.au